

### Solution 1

(a) $m\ddot{X}_n = S(X_{n+1} - X_n) - S(X_n - X_{n-1})$ .	0.7
(b) Let $X_n = A \sin nka \cos(\omega t + \alpha)$ , which has a harmonic time dependence. By analogy with the spring, the acceleration is $\ddot{X}_n = -\omega^2 X_n$ .	
Substitute into (a): $-mA\omega^2 \sin nka = AS \{\sin(n+1)ka - 2 \sin nka + \sin(n-1)ka\}$	
$= -4SA \sin nka \sin^2 ka$ .	0.6
Hence $\omega^2 = (4S/m) \sin^2 ka$ .	0.2
To determine the allowed values of $k$ , use the boundary condition $\sin(N+1)ka = \sin kL = 0$ .	0.7
The allowed wave numbers are given by $kL = \pi, 2\pi, 3\pi, \dots, N\pi$ ( $N$ in all),	0.3
and their corresponding frequencies can be computed from $\omega = \omega_0 \sin ka$ ,	
in which $\omega_{\max} = \omega_0 = 2(S/m)^{1/2}$ is the maximum allowed frequency.	0.4
(c) $\langle E(\omega) \rangle = \frac{\sum_{p=0}^{\infty} p\hbar\omega P_p(\omega)}{\sum_{p=0}^{\infty} P_p(\omega)}$	
First method: $\frac{\sum_{n=0}^{\infty} n\hbar\omega e^{-n\hbar\omega/k_B T}}{\sum_{n=0}^{\infty} e^{-n\hbar\omega/k_B T}} = k_B T^2 \frac{\partial}{\partial T} \ln \sum_{n=0}^{\infty} e^{-n\hbar\omega/k_B T}$	1.5
The sum is a geometric series and is $\{1 - e^{-\hbar\omega/k_B T}\}^{-1}$	0.5
We find $\langle E(\omega) \rangle = \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$ .	
<i>Alternatively:</i> denominator is a geometric series = $\{1 - e^{-\hbar\omega/k_B T}\}^{-1}$	(0.5)
Numerator is $k_B T^2 (d/dT)$ (denominator) = $e^{-\hbar\omega/k_B T} \{1 - e^{-\hbar\omega/k_B T}\}^{-2}$ and result follows.	(1.5)

*A non-calculus method:*

Let  $D = 1 + e^{-x} + e^{-2x} + e^{-3x} + \dots$ , where  $x = \hbar\omega/k_B T$ . This is a geometric series and equals  $D = 1/(1 - e^{-x})$ . Let  $N = e^{-x} + 2e^{-2x} + 3e^{-3x} + \dots$ . The result we want is  $N/D$ . Observe

$$\begin{aligned} D - 1 &= e^{-x} + e^{-2x} + e^{-3x} + e^{-4x} + e^{-5x} + \dots \\ (D - 1)e^{-x} &= e^{-2x} + e^{-3x} + e^{-4x} + e^{-5x} + \dots \\ (D - 1)e^{-2x} &= e^{-3x} + e^{-4x} + e^{-5x} + \dots \end{aligned}$$

Hence  $N = (D - 1)D$  or  $N/D = D - 1 = \frac{e^{-x}}{1 - e^{-x}} = \frac{1}{e^x - 1}$ .

(2.0)

(d) From part (b), the allowed  $k$  values are  $\pi/L, 2\pi/L, \dots, N\pi/L$ .

Hence the spacing between allowed  $k$  values is  $\pi/L$ , so there are  $(L/\pi)\Delta k$  allowed modes in the wave-number interval  $\Delta k$  (assuming  $\Delta k \gg \pi/L$ ).

1.0

(e) Since the allowed  $k$  are  $\pi/L, \dots, N\pi/L$ , there are  $N$  modes.

0.5

Follow the problem:

$d\omega/dk = a\omega_0 \cos ka$  from part (a) & (b)

$$= \frac{1}{2} a \sqrt{\omega_{\max}^2 - \omega^2}, \quad \omega_{\max} = \omega_0. \text{ This second form is more convenient for integration.}$$

The number of modes  $dn$  in the interval  $d\omega$  is

$$\begin{aligned} dn &= (L/\pi)\Delta k = (L/\pi) (dk/d\omega) d\omega \\ &= (L/\pi) \{a\omega_0 \cos ka\}^{-1} d\omega \end{aligned}$$

0.5 for eitl

$$\begin{aligned} &= \frac{L}{\pi} \frac{2}{a} \frac{1}{\sqrt{\omega_{\max}^2 - \omega^2}} d\omega \\ &= \frac{2(N+1)}{\pi} \frac{1}{\sqrt{\omega_{\max}^2 - \omega^2}} d\omega \end{aligned}$$

This part is necessary for  $E_T$  below,

but not for number of modes

Total number of modes =  $\int dn = \int_0^{\omega_{\max}} \frac{2(N+1)}{\pi} \frac{d\omega}{\sqrt{\omega_{\max}^2 - \omega^2}} = N + 1 \approx N$  for large  $N$ .

(0.5)

Total crystal energy from (c) and  $dn$  of part (e) is given by

$$E_T = \frac{2N}{\pi} \int_0^{\omega_{\max}} \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1} \frac{d\omega}{\sqrt{\omega_{\max}^2 - \omega^2}}.$$

0.7

(f) Observe first from the last formula that  $E_T$  increases monotonically with temperature since

$\{e^{\hbar\omega/k_B T} - 1\}^{-1}$  is increasing with  $T$ .

---

0.2

When  $T \rightarrow 0$ , the term  $-1$  in the last result may be neglected in the denominator so

---

0.2

$$E_T \approx_{T \rightarrow 0} \frac{2N}{\pi} \int \hbar\omega e^{-\hbar\omega/k_B T} \frac{1}{\sqrt{\omega_{\max}^2 - \omega^2}} d\omega$$

$$= \frac{2N}{\hbar\pi\omega_{\max}} (k_B T)^2 \int_0^{\infty} \frac{x e^{-x}}{\sqrt{1 - (k_B T x / \hbar\omega_{\max})^2}} dx$$

---

0.3

---

0.2

which is quadratic in  $T$  (denominator in integral is effectively unity) hence  $C_V$  is linear in  $T$  near absolute zero.

---

0.2

Alternatively, if the summation is retained, we have

$$E_T = \frac{2N}{\pi} \sum_{\omega} \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1} \frac{\Delta\omega}{\sqrt{\omega_{\max}^2 - \omega^2}} \rightarrow_{T \rightarrow 0} \frac{2N}{\pi} \sum_{\omega} \hbar\omega e^{-\hbar\omega/k_B T} \frac{\Delta\omega}{\sqrt{\omega_{\max}^2 - \omega^2}}$$

$$= \frac{2N}{\pi} \frac{(k_B T)^2}{\hbar\omega} \sum_y e^{-y} y \Delta y$$

---

(0.5)

When  $T \rightarrow \infty$ , use  $e^x \approx 1 + x$  in the denominator,

---

0.2

$$E_T \approx_{T \rightarrow \infty} \frac{2N}{\pi} \int_0^{\omega_{\max}} \frac{\hbar\omega}{\hbar\omega/k_B T} \frac{1}{\sqrt{\omega_{\max}^2 - \omega^2}} d\omega = \frac{2N}{\pi} k_B T \frac{\pi}{2},$$

---

0.1

which is linear; hence  $C_V \rightarrow Nk_B = R$ , the universal gas constant. This is the Dulong-Petit rule.

Alternatively, if the summation is retained, write denominator as  $e^{\hbar\omega/k_B T} - 1 \approx \hbar\omega/k_B T$  and

---


(0.2)

$$E_T \rightarrow_{T \rightarrow \infty} \frac{2N}{\pi} k_B T \sum_{\omega} \frac{\Delta\omega}{\sqrt{\omega_{\max}^2 - \omega^2}} \text{ which is linear in } T, \text{ so } C_V \text{ is constant.}$$

Sketch of  $C_V$  versus  $T$ :

---

0.5



**Answer sheet: Question 1**

(a) Equation of motion of the  $n^{\text{th}}$  mass is:

$$m\ddot{X}_n = S(X_{n+1} - X_n) - S(X_n - X_{n-1}).$$

(b) Angular frequencies  $\omega$  of the chain's vibration modes are given by the equation:

$$\omega^2 = (4S/m) \sin^2 ka.$$

Maximum value of  $\omega$  is:  $\omega_{\text{max}} = \omega_0 = 2(S/m)^{1/2}a$

The allowed values of the wave number  $k$  are given by:

$$\pi/L, 2\pi/L, \dots, N\pi/L.$$

How many such values of  $k$  are there?  $N$

(f) The average energy per frequency mode  $\omega$  of the crystal is given by:

$$\langle E(\omega) \rangle = \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$$

(g) There are how many allowed modes in a wave number interval  $\Delta k$ ?


$$(L/\pi)\Delta k.$$

(e) The total number of modes in the lattice is:  $N$

Total energy  $E_T$  of crystal is given by the formula:

$$E_T = \frac{2N}{\pi} \int_0^{\omega_{\max}} \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1} \frac{d\omega}{\sqrt{\omega_{\max}^2 - \omega^2}}.$$

(h) A sketch (graph) of  $C_V$  versus absolute temperature  $T$  is shown below.



For  $T \ll 1$ ,  $C_V$  displays the following behaviour:  $C_V$  is linear in  $T$ .

As  $T \rightarrow \infty$ ,  $C_V$  displays the following behaviour:  $C_V \rightarrow Nk_B = R$ , the universal gas constant.